

Submission on the proposed National Policy Statement for Indigenous Biodiversity

Professor David Norton
Te Kura Ngahere | School of Forestry
University of Canterbury, Christchurch
david.norton@canterbury.ac.nz or 027-201-7794

9 March 2020

While I accept that a consistent approach towards managing indigenous biodiversity across New Zealand is a desirable outcome, I have significant doubts about the ability of the proposed NPSIB to deliver the outcomes that its proponents suggest it can. My concerns centre primarily around (1) the way it is proposed that significant indigenous biodiversity will be assessed and (2) the reliance on a rules-based approach to managing significant indigenous biodiversity without the provision of genuine support (incentives, extension etc) for landowners who have significant indigenous biodiversity.

I believe that I am uniquely placed to make these comments. I have over 30 years experience undertaking research on indigenous biodiversity in rural New Zealand, including a 2013 book¹ and numerous presentations and publications on the ways to achieve win-win outcomes for biodiversity and farming. I have worked closely with the farming sector in developing planning approaches to incorporating biodiversity into farm management and have a good sense of how best to incentivise farmers to manage indigenous biodiversity. At the same time, I have researched and published widely across a range of ecological and conservation issues in New Zealand such as threatened plant conservation and biodiversity offsetting, including co-authoring the paper that established the system we use in New Zealand today for classifying the conservation status of all flora and fauna. I would suggest that there are few other ecologists in New Zealand with the type of experience that I have.

And I wanted to emphasise at the outset that what has motivated me for my 40-odd years as a university researcher and teacher has been to find ways to enhance New Zealand's unique indigenous biodiversity. But in order to achieve this, I have always recognised that we also require a vibrant economy to pay for conservation of which primary production has been and will continue to be a critical component.

I have chosen not to comment in detail on the proposed NPSIB as I believe that the overall proposal is flawed and should be redrafted to address these higher-level concerns. I raised these concerns with the Biodiversity Collaborative Group in a presentation I made to them on 5 July 2018 but the issues that I raised seem to have been ignored in the drafting of the NPSIB which I find very disappointing. I am strongly of the opinion that the proposed NPSIB will result in a net loss of indigenous biodiversity, not because farmers will deliberately seek to destroy it (although I suspect some are doing this right now), but because the rules-based approach proposed in the NPSIB will disenfranchise most farmers from wanting to look after indigenous biodiversity and it will decline through benign neglect. I think that this is a massive tragedy and a real lost opportunity for New Zealand conservation and one that we will come to regret in the future.

¹ Norton D, Reid N 2013. Nature and farming. Sustaining native biodiversity in agricultural landscapes. Melbourne, CSIRO Publishing.

Identification of significance: I believe that the criteria proposed in the NPSIB are too broad and will include virtually all areas with indigenous vegetation or that support indigenous fauna in New Zealand. While conservation of indigenous biodiversity is of fundamental importance, an approach that makes all indigenous biodiversity including substantial areas of regenerating woody vegetation dominated by species like kānuka and matagouri significant fails to recognise that some areas of indigenous biodiversity are much more important/significant than others and that their protection and enhancement is a much higher priority than other areas, especially as resources for their management is limited.

Matagouri is a good example of this. The presence of matagouri on a farm triggers the proposed significance criteria as matagouri is ranked nationally as 'At Risk – Declining'. However, in eastern South Island hill and high country matagouri responds vigorously to fertiliser application and can be widespread across hill sides that have a long history of fertiliser application and pastoral farming. These sites typically have few other indigenous species and the matagouri shrublands present usually have a dense ground cover of exotic grasses and herbs. Most farmers want to retain a degree of matagouri cover (it provides important shelter for lambing and in droughts, grass growth is usually better in its shade), but also need to be able to control matagouri so it does not become too dense. Defining these areas as significant under the proposed NPSIB criteria makes it very difficult to convince a farmer why another area of shrubland dominated by genuinely threatened species such as *Leonohebe cupressoides* or *Olearia fimbriata* is also significant and actually requires proactive protective management.

Similar arguments can be made for kānuka and mānuka. While important early successional species in the establishment of native forest on farmland, kānuka and mānuka have established widely across rural New Zealand on previously pasture dominated sites. While the NPSIB does include some exemptions around kānuka and mānuka, the way the criteria are worded it seems to me highly likely that almost all areas of kānuka and mānuka will be deemed as significant by territorial local authorities (TLAs) under the proposed NPSIB criteria. Again, convincing a farmer that a stand of kānuka or mānuka that has clearly regenerated on a previously pasture-dominated hill side has the same value as a cutover remnant of old growth podocarp-tawa forest is going to be very difficult.

My point here is that we do need to have significance criteria, but as we argued in a 2004 paper (Norton & Lindsay 2004²) these need to focus on the most important sites, the remnants of indigenous biodiversity that we need to protect and enhance as a priority. Then we can use other methods to achieve protection and management of other areas. Making the criteria too broad means that it will be much harder to help landowners and the general public to understand why some sites are more important/significant than others, and therefore why we need to prioritise what are already scarce conservation resources (especially at a TLA level) to focus on some sites over others. Broad criteria will also make TLAs the 'piggy in the middle' in what are likely to be protracted arguments between landowners and those advocating for this broad definition of significance.

Implementing biodiversity conservation on private land: I believe strongly that we need to clearly distinguish between the objective ecological process of significance assessment

² Norton DA, Roper-Lindsay J 2008. Assessing significance for biodiversity conservation on private land in New Zealand. *New Zealand Journal of Ecology* 28: 295-305.

(preferably using a more focused approach than that proposed in the NPSIB) and the more subjective processes of providing for the protection of the values once identified. Simply calling an area significant will **not** contribute in any way towards sustaining and preferably enhancing the values the site contains. It is totally naïve to think otherwise, yet this is essentially what the proposed NPSIB does and I see this as a real failing of the proposed NPSIB. I wrote an article in 1988³ that argued just this – history suggests that we are slow at learning! The approach advocated in the NPSIB is one that is largely rules-based and while some degree of grandparenting is provided for (Section 3.12), I do not believe that the rules-based approach is appropriate or will deliver the outcomes that it hopes to deliver. In particular, I find it really concerning that there is nothing in the NPSIB about systems to incentivise and support landowners manage indigenous biodiversity on their land. However, I do acknowledge that in the Discussion Document there is discussion about guidance and support for implementing the proposed NPSIB (Section E7). But these suggestions are outside the proposed NPS itself and therefore there are no guarantees that they will be enacted or given effect to by any level of Government. This is a major failure of the proposed NPSIB.

We have recently published a paper (Norton et al 2020⁴) that argues that if we are to achieve the desired outcomes for indigenous biodiversity in rural New Zealand, landowners need to be incentivised and rewarded for good biodiversity management (not penalised through a rules-based system). To bring integrity and objective support to such an incentive-based approach to biodiversity conservation in rural New Zealand, we suggest that farmers need to adopt environmental management planning that is supported by good biodiversity extension resources and with a verification system that shows farmers are doing what they say they are doing. This approach requires **trust** and **partnership** among all players in agroecosystems – farmers, government, food and fibre processors, scientists, conservationists, NGOs, and the wider New Zealand population. I do not believe that the proposed NPSIB encourages or supports such trust or partnership. Our paper concludes by suggesting that if we change the way we think about how farming and biodiversity interact (and move away from a rules-based approach), then we will achieve substantial biodiversity gains across the 50% of New Zealand under pastoral farming. As I suggested at the start of this submission, a rules-based approach will not result in biodiversity gains and will in fact result in biodiversity loss.

I will not repeat the content of this 2020 paper but append it to this document as a core part of my submission on the NPSIB. But I would like to reiterate the points made above that in order to achieve any meaningful gains in indigenous biodiversity across rural New Zealand, it is essential that there is a meaningful system of incentives and support for rural landowners. That we have a proposed NPSIB signals that indigenous biodiversity is a public good and therefore must be supported by public funding. The core of this support must include a decent extension system to support farmers and meaningful financial incentives for on-the-ground biodiversity management (including compensation).

I understand that there will not be any public hearings associated with further development of the proposed NPSIB but I am very happy to provide further input and clarification around my comments if required.

³ Norton DA 1988. Managing for the long term. *Forest and Bird* 19(2): 32–34.

⁴ Norton DA, Suryaningrum F, Buckley HL², Case BS, Cochrane CH, Forbes AS, Harcombe M 2020. Achieving win-win outcomes for pastoral farming and biodiversity conservation in New Zealand. *New Zealand Journal of Ecology* 44(2): in press.

New Zealand Journal of Ecology 44(2), 2020, in press.

Achieving win-win outcomes for pastoral farming and biodiversity conservation in New Zealand

David A. Norton^{1*}, Febyana Suryaningrum², Hannah L. Buckley², Brad S. Case², C. Hamish Cochrane³, Adam S. Forbes¹ and Matt Harcombe⁴

¹Te Kura Ngahere/School of Forestry, University of Canterbury, Christchurch 8014, New Zealand

²School of Science, Auckland University of Technology, Auckland 1142, New Zealand

³Capra hircus Consulting, C/- 3 Picton Ave, Christchurch 8011, New Zealand

⁴3 Conway Street, Dunedin 9014, New Zealand

*Author for correspondence (Email: david.norton@canterbury.ac.nz)

Published online:

Abstract: Pastoral farming is the dominant land use in New Zealand today and is under considerable domestic social and political pressure to reduce its environmental footprint. In this article, we explore options to enhance native biodiversity conservation within New Zealand pastoral systems. We argue that there is strong synergistic interdependence between biodiversity conservation and pastoral farming and suggest that it is possible to have win-win outcomes for both. Landowners need to be incentivised and rewarded for good biodiversity management, rather than relying on a strict rules-based approach. To bring integrity and objective support to this incentive-based approach, farmers need to adopt environmental management planning that is supported by good biodiversity extension resources. Alongside this, a verification system is required that shows farmers are doing what they say they are doing and reflects agreed management targets for biodiversity. This approach requires trust and partnership among all players in agroecosystems – farmers, government, food and fibre processors, scientists, conservationists, NGOs, and the wider New Zealand population. We suggest that if we change the way we think about how farming and biodiversity interact, then we will achieve substantial biodiversity gains across the 50% of New Zealand under pastoral farming. This then brings integrity to the existing and expanding market story for pastoral farming and creates a stronger connection between all New Zealanders and the farming sector. Advancing our thinking in this way will enable New Zealand to maintain a premium for our farming products internationally while supporting conservation of our native biodiversity.

Keywords: agroecosystems, biodiversity, conservation, ecosystem services, environmental verification, incentives, pastoral farming, policy

Introduction

Pastoral farming is the dominant land use in New Zealand, occupying at least half of the land area (40% sheep and beef, and 10% dairy; Norton & Pannell 2018), and is a critical part of the New Zealand economy contributing 36% and 38% of export earnings (including international tourism) in 2017 and 2018 respectively (Stats NZ pers. comm. 2019). Pastoral farming occurs primarily at lower elevations where loss of original habitat has been greatest (Ewers et al. 2006; Perry et al. 2014) and where there is little public conservation land,

although significant remnants of native vegetation still remain (Norton & Pannell 2018). Before humans reached New Zealand (1230–1280 AD; Wilmshurst et al. 2011), the areas where pastoral farming currently occurs were dominated by diverse, mostly forested ecosystems (Leathwick 2001), although grasslands and shrublands were present in some parts of the eastern South Island high country (McGlone 2000), and wetlands, although small, occur throughout. The temperate climates, reasonable rainfall and fertile soils that favoured diverse native forests have made these areas well suited to pastoral farming (Leathwick et al. 2001).

Pastoral farming is under considerable domestic social and political pressure to reduce its environmental footprint. Substantial concerns have been raised about the impact of farming systems (especially high intensity dairy farming) on water quality (Foote et al. 2015) and the contribution of grazing animals to greenhouse gas emissions, especially biogenic methane (PCE 2019; MfE 2019a). Internationally, the agricultural sector, especially the animal-based farming sector, is also under increasing pressure. Concerns around the contribution of animal farming to greenhouse gas emissions, biodiversity loss and animal welfare and health are all increasing (Newbold et al. 2015; Poore & Nemecek 2018, Almiron & Tafalla 2019), with calls for a significant shift away from current consumption of red meat and dairy products towards synthetic and plant-based alternatives.

Agriculture's impact on biodiversity has been well documented both globally (e.g. Tilman 1999, Potts et al. 2010) and locally (e.g. Dale et al. 1994, Butler et al. 2010). In the 2019 IPCC Special Report on Climate Change and Land (IPCC 2019a), expansion of agriculture and forestry as a result of Earth's growing human population has contributed to 23% of the total anthropogenic greenhouse gas emissions, as well as increasing the loss of natural ecosystems, contributing to declining biodiversity. It seems clear that as the global human population continues to grow, food production and distribution will need to become more efficient. Notwithstanding this, pressures from farming on biodiversity are likely to continue (Crist et al. 2017; Balmford et al. 2019). Land sparing and land sharing have been proposed as two ways of conceptualising how we might manage biodiversity within agroecosystems, although the consensus seems to be that both are likely to be important, especially because land set aside for biodiversity conservation under the land sparing model is vulnerable to future impacts (Fischer et al. 2014). While land sparing and land sharing can be seen as opposite ends of a continuum of approaches to biodiversity management within agroecosystems, the relative importance of either is strongly influenced by both the social-ecological history of the area under consideration and the spatial scale at which landscape elements are being considered (Grass et al. 2019).

In this article, we explore options to enhance native biodiversity conservation within New Zealand pastoral agroecosystems while also optimising primary production. We believe that there is strong synergistic interdependence between biodiversity conservation and pastoral farming that is not well recognised, and that it is possible to have win-win outcomes for both in New Zealand. However, to do this we suggest the need for a fundamental shift both in our farming systems and in the way we implement and support biodiversity management and enhancement. This shift needs to be away from traditional thinking where biodiversity conservation or pastoral farming are viewed as a dominant land management objective on their own to one where they are spatially integrated and complement each other. We focus here primarily on forested ecosystems because, with the exception of parts of the eastern South Island high country, forest was the predominant pre-European vegetation cover in the areas where pastoral farming now occurs.

Biodiversity and pastoral farming – a brief overview

Biodiversity in New Zealand pastoral systems

The composition, structure and spatial arrangement of native biodiversity in those parts of New Zealand where pastoral agroecosystems dominate is different to those in areas with extensive public conservation lands. With some exceptions, native habitats tend to be small, isolated, and modified to varying degrees by grazing (domestic and feral), historic logging, plant invasions and edge effects resulting in changes to their composition and structure (Timmins & Williams 1991; Burns et al. 2000; Smale et al. 2008; Burns et al. 2011; Ruffell & Didham 2017). While some of these native habitats are protected through covenants or as part of the public conservation estate, most do not have any formal protection (Norton & Pannell 2018). The fragments of native habitat that occur comprise both remnants of the original forests and regenerating vegetation that has established on sites that were previously farmed. This type of woody native vegetation can range from a continuous native canopy to scattered trees through a paddock.

Notwithstanding its fragmented and isolated nature, a surprisingly large amount of native biodiversity remains across pastoral landscapes in rural New Zealand, including many nationally and regionally rare plant and animal species (e.g. MacLeod et al. 2008; de Lange et al. 2010; Pawson et al. 2010). While species requiring large areas of intact habitat such as kōkako *Callaeas cinerea* and kākā *Nestor meridionalis* are usually absent, other nationally rare species still sporadically persist (e.g. pōpokatea *Mohoua albicilla* and brown kiwi *Apteryx mantelli* in the North Island). As is the case on public conservation land, many native plants, birds, lizards and invertebrates are in decline in pastoral agroecosystems because of the ongoing effects of predation pressure, as well as the legacy effects of fragmentation (loss of resources and landscape connectivity). Nonetheless, the remaining native biodiversity that is present in pastoral landscapes is critically important across extensive areas of lowland New Zealand because these remnants are often all that remains of the original pre-human ecosystems (Norton & Pannell 2018).

Pastoral farming systems

New Zealand farming systems are still largely grass-based and display a continuum from relatively extensive, low-input systems to high-input intensive farming systems. Intensification has been characterised by increases in stocking rate, fertiliser use, irrigation area, cultivation techniques and extent, and smaller paddock sizes (MacLeod & Moller 2006; Didham et al. 2015). Much of this has been driven by the expansion of dairy farming, often replacing sheep and beef farming. Irrigation is often seen as a key indicator of intensification, although the increasing area of irrigated land is not just a result of dairy expansion, with significant irrigation increases occurring in other sectors including sheep and beef farming (17% irrigated area), arable farming (13%), and horticulture including viticulture (11%) (see: www.stats.govt.nz/indicators/irrigated-land). Intensification across a range of landscapes, but especially associated with dairy farming, has resulted in reductions in stream flows, increases in nutrient discharges into streams and groundwater, and eutrophication of wetlands (Foote et al. 2015; Ramezani et al. 2016). However, in some farming systems intensification has actually improved environmental outcomes. For example, the shift from flood irrigation (wild flood and border dyke) to pivot irrigation on some pastoral farms has led to improved environmental outcomes for these systems due to less overland flow resulting in a decrease in leaching losses and nutrient concentrations in receiving water bodies (McDowell 2017).

Plantation forestry, primarily with radiata pine (*Pinus radiata*), has also expanded over the last few decades in traditional sheep and beef farming areas (Wallace 2019). Government policy initiatives (e.g. Erosion Control Funding Programme in Gisborne District and the Hill Country Erosion Fund more generally) have facilitated this expansion (www.teururakau.govt.nz/funding-and-programmes/environment-and-natural-resources/erosion-control-funding-programme/). Many farmers have also been actively establishing trees, woodlots and farm forestry on their properties for erosion control, to earn carbon credits through the Emissions Trading Scheme (ETS), and as a future income source for retirement and succession planning (Kennet et al. 2010, Hutching 2019). With an increasing focus on the effects of climate change, the One Billion Trees Programme (Te Uru Rākau 2018), the New Zealand Government's target of being carbon-zero by 2050, and a predicted rising value of carbon credits (www.carbonnews.co.nz), trees may become a more attractive investment for some sheep and beef farms than traditional farming (Timar 2016). These more recent policy initiatives have triggered another shift from sheep and beef farming to exotic plantation forestry for carbon credits with suggestions that as much as 30 000 ha in the eastern North Island has been converted in 2018–19 (Wallace 2019).

Regulatory environment

There is increasing pressure on farmers from markets, different levels of government, environmental NGOs and society generally, to be more environmentally responsible (Gabzdylova et al. 2009; Mitchell 2017; Diprose 2018; MfE 2019a). District and regional councils are increasingly regulating farmers in terms of how they interact with biodiversity, primarily through the use of planning tools (Brown 2016), including restrictions on vegetation clearance, pastoral intensification, and riparian management. Several district and regional councils now require farmers to have farm environment plans that include biodiversity in order to obtain or retain resource consents for a variety of on-farm management activities (e.g. irrigation, fertiliser application, vegetation clearance; Blaschke & Ngapo 2003). Central Government has recently released a draft National Policy Statement for Indigenous Biodiversity which proposes even stricter requirements on farmers in terms of indigenous biodiversity (MfE 2019b).

At the same time, there is increasing recognition within the farming sector of the need to prove its environmental credentials as part of its social licence to operate (e.g. Dairy Tomorrow 2017, B+LNZ 2018a). Many individual farmers are responding positively through retiring bush and wetland areas from farming and in some cases covenanting them. However, covenants (mainly through the QEII National Trust) only account for 3% of the total area of native vegetation on sheep and beef farms (Norton & Pannell 2018). Some farmers are reluctant to covenant or otherwise formally set aside areas on their farms, fearing that this might compromise future options especially if regulation becomes more restrictive. This fear of regulation also has the potential to lead to perverse outcomes such as farmers being prepared to accept fines for vegetation clearance as a better outcome than what might be perceived as a likely expensive and drawn-out resource consent process. Notwithstanding this, the uptake of farm management plans (also called farm or land environment plans) has been increasing (Maseyk et al. 2019) and Beef + Lamb New Zealand have called for all sheep and beef farmers to have next generation land environment plans for their farms that include consideration of water, soil, carbon and biodiversity as the four pillars for sustainable farming (B+LNZ 2018a).

Government has, and continues to, put in place policies encouraging tree planting, including native trees, in agroecosystems. The Emissions Trading Scheme and Permanent Forest Sink

Initiative together with various afforestation grants work to discourage deforestation and encourage reforestation by providing an avenue for the creation of permanent forests (Leining & Kerr 2018). The ETS was set up to reduce deforestation in old forest (pre-1990 exotic forest) and incentivise new forest (post-1989 native and exotic forest). While the ETS and voluntary carbon-trading systems incentivise landowners to establish native and exotic trees, or promote natural regeneration, the direct benefits to biodiversity through reforestation have been limited because significantly more carbon credits can be earned through exotic plantations as low carbon prices generally make native afforestation uneconomic. Furthermore, under the ETS, a carbon-credit earning forest can still be harvested, as long as it is replanted, providing another income source from timber yield (Leining & Kerr 2018) which again does not promote biodiversity conservation. The One Billion Trees programme (Te Uru Rākau 2018) offers further incentives for farmers to integrate trees into their landscape by helping fund land retirement, fencing and the planting of native species. With the recently legislated Zero Carbon Act 2019 and the One Billion Trees programme, coupled with rising carbon prices (www.carbonnews.co.nz), there is now the very real potential for land owners to take advantage to optimise income through carbon sequestration by planting or enhancing existing stands with native woody species providing significant co-benefits and opportunities for enhancing biodiversity in working pastoral landscapes.

Can we better integrate biodiversity conservation and pastoral farming?

The answer is yes, but to do this we suggest a rethink of the approach towards both biodiversity conservation and pastoral farming in rural landscapes is required; each depends on the other, addressing one in isolation will not result in a sustainable future for either (Norton & Reid 2013). What is required is for New Zealanders to engage in social and ecological discussions about the future of these systems; we need to be asking what will make these systems sustainable in 2050 and beyond? We believe that these discussions need to consider several propositions in order to obtain sustainable win-win outcomes for biodiversity conservation and pastoral farming (Table 1).

Table 1. Key propositions underpinning biodiversity conservation in pastoral systems.

Proposition	Explanation
Change the approach to biodiversity conservation	Biodiversity conservation in pastoral agroecosystems will be different to that on public conservation land in terms of how it is done and the outcomes achieved. However, it will also be complementary, and contribute significantly, to regional and national biodiversity conservation by helping to reverse the declines in native species across all of New Zealand.
Prove biodiversity conservation is occurring in agroecosystems	Evidence of active biodiversity management and resultant outcomes is essential for pastoral farming as part of its social licence to operate domestically and to market value-added products globally.
Show that biodiversity benefits farming	Biodiversity enhances ‘multifunctionality’ in agroecosystems and landscapes (e.g. cultural benefits, carbon sequestration, alternative income sources and reversing environmental damage such as water quality and quantity) and leads to better on-farm human wellbeing (Maseyk et al. 2017) and animal health.
Farming is economically sustainable	Economically-sustainable pastoral farming businesses are essential to fund biodiversity conservation work across pastoral agroecosystems, both through individual farms having the financial freedom to do their own biodiversity work and for government to collect sufficient rates and tax income to fund government’s biodiversity work (Norton and Reid 2013).
Support farmers to achieve biodiversity goals	Good management planning, extension and demonstration services, verification and incentive/reward systems to support farmers are essential for good biodiversity outcomes in agroecosystems.

In developing a new future for pastoral farming and biodiversity conservation, we need to address some fundamental issues:

- (1) The first is that conservation groups and the wider New Zealand public need to trust that, if properly incentivised, valued and resourced, farmers are the best stewards of native biodiversity on their land (Norton & Reid 2013). The Department of Conservation does not have the resources to manage the public conservation estate let alone work with the myriad of small habitat patches dotted across rural New Zealand. District and regional councils also lack the resources, expertise, trust and mandate to do this. A recent phone survey of 500 sheep and beef farmers showed that nearly 80% felt that protecting and managing native biodiversity on their farms was important (Maseyk et al. unpubl. data).
- (2) At the same time, it is important that farmers recognise that there are many groups and individuals outside their farms who have an interest in and expertise about biodiversity across pastoral agroecosystems and want to help (directly or indirectly) achieve positive biodiversity outcomes on farms. Trust needs to work both ways.
- (3) To make biodiversity conservation economically viable in agroecosystems we need to focus on producing value-added products that can be differentiated, marketed, verified and sold based on their method of production and on their environmental/biodiversity story. The current Beef+Lamb NZ campaign “Taste Pure Nature” is a good example of this.
- (4) In developing a new future for pastoral farming and biodiversity conservation, contributions from all the different groups involved (private sector, consumers, land managers, local communities, Māori, policy makers) to create enabling conditions for this and to scale up to regional and international markets (IPCC 2019b.).
- (5) Finally, we need to enhance the current and proposed (through the National Policy Statement for Indigenous Biodiversity) regulatory approach to managing biodiversity in agroecosystems taken in District and Regional Plans, by incentivising and rewarding farmers for their biodiversity conservation actions. Such changes to the current approach need to be based on trust, partnership and collaboration.

In managing native biodiversity in agroecosystems, it is important to recognise that the amount of native biodiversity on individual farms will vary, but it is difficult to conceive any pastoral farms that do not have some native biodiversity, even if only visits by mobile species such as pūtangitangi *Tadorna variegata* and korimako *Anthornis melanura*, or riparian restoration plantings. All pastoral farmers need to understand and appreciate the value of whatever native biodiversity they have, including its significance in terms of criteria defined in district and regional plans, and incorporate biodiversity into their farm planning systems. To do this, biodiversity management should be seen as a collaboration that involves everyone from enviroschools and kids, to scientists and conservation groups, to local and central Government, supporting farmers in appropriate ways to manage biodiversity, while also recognising that farmers need to be operating profitable businesses that benefit not only farming families and their immediate communities, but all of New Zealand.

How do we achieve win-win outcomes for native biodiversity and pastoral production in agroecosystems?

To achieve win-win outcomes we need to better target biodiversity management on farms, do our farming smarter and provide genuine support for farmers in implementing biodiversity management on their farms.

Targeted biodiversity management

Before we commence on the how, let's be clear about what we are collectively trying to achieve. Protection of all that remains of our original old-growth forests and wetlands, and the extensive areas of regenerating native forest is essential (Norton et al. 2018). The sustainability of remnants of the original old growth forests is critical even when they might have been impacted in some manner through historic logging, grazing animals or the effects of adjacent agricultural activities (Smale et al. 2008; Burns et al. 2011; Didham et al. 2015). These remnants are the direct connections with the past and the source of propagules (plant, animal, fungal) for the future. Regenerating forest represents systems that are already developing towards a more mature phase and, through appropriate management, are a low-cost way to enhance native biodiversity at landscape scales. Simply protecting these areas legally or even just fencing them off is not, however, sufficient to guarantee their viability (Norton 1988). What is required is the implementation of management practices in remnants and regenerating vegetation to enhance them so that native biodiversity flourishes (e.g. Dodd et al. 2011; Ruffell & Didham 2017). This need not exclude economic uses (shelter, timber, grazing, honey etc) depending on the local situation and the values present.

Restoration of native habitat on farms is also important, both for ensuring that key resources for native fauna are present, such as year-round food supplies and nesting opportunities, and for enhancing connectivity across landscapes (Richard & Armstrong 2010). Restoration can involve a mix of strategic planting and seeding, facilitated natural regeneration, and enhancement of existing areas of degraded and regenerating forest through enrichment (Norton et al. 2018; Forbes et al. 2020). Exotic habitats, such as farm woodlots and homestead gardens, can also contribute to enhancing connectivity and increasing resource availability for native fauna (Norton 1998; Campbell et al. 2008; McArthur et al. 2019). Management of remnants, regenerating vegetation and restoration requires plant and animal pest control, especially of herbivores and omnivores (deer, goat *Capra hircus*, pig *Sus scrofa*, possum *Trichosurus vulpecula*), as well as consideration of interactions between different land uses (e.g. fertiliser drift). However, addressing predation (carnivore) pressure alone without enhancing resources and increasing connectivity is insufficient for biodiversity conservation and will not result in enhancement of all native biodiversity in agroecosystems, especially of mobile species such as birds (Pannell et al. unpubl. data).

Biodiversity conservation activities on farms might occur through land sparing (taking land out of production for restoration), but it will also occur within the farmed landscape itself. Enhancing and conserving biodiversity might involve using a different mix of species for woodlots, erosion plantings and shelterbelts, retaining and planting scattered native trees in paddocks (Manning et al. 2006; Fukuda et al. 2011), shrub retention (e.g. matagouri *Discaria toumatou*), ensuring that year-round food supplies are available for native birds and invertebrates (including exotic tree sources), and taking advantage of recent advances in precision agriculture to minimise farming impacts on areas of native habitat. Appropriate economic use of remnants and regenerating vegetation may also be appropriate so long as the core values present are not compromised.

The key is taking a landscape-scale perspective on farming and biodiversity conservation, making sure that farming is undertaken in a manner that is friendly to biodiversity (land sharing), and recognising that much of this native biodiversity can directly benefit farming through shelter, shade, soil retention and nutrient management (Balmford et al. 2019; Case et al. unpubl. data) and that farming can benefit biodiversity (Norton & Reid 2013). While there are still gaps in our knowledge of native biodiversity in agroecosystems such as

dispersal patterns for mobile native species (Norton 2001; MacLeod et al; 2008; Pannell et al. in submission) we are suggesting the critical factor is to be able to design our agroecosystems using an approach that optimises both biodiversity management and pastoral farming at the landscape scale.

Smarter farming

Underpinning our biodiversity management, continuing to get smarter with how we farm is equally as important. This needs to include matching the right animals, grazing practices and forage plants to the right parts of the farm. We can continue to improve how we do this through better using and mapping land use capability and developing specific actions for identified land management units (Dominati et al. 2016). This might include adapting animal husbandry to different areas of the farm (e.g. different forms of rotational grazing), picking forage plants that best suit the environment (especially with climate change), reducing nutrient losses by targeting animal type and fertiliser use, thinking more about soil management, retiring difficult-to-farm areas or making the most of technology to optimise production in each of those land management units (e.g. precision agriculture). New Zealand pastoral farming has already done much in this space, with sheep and beef farmers for example producing the same amount of product with fewer animals today than in the past (B+LNZ 2018b). Adopting or enhancing some of the ideas of regenerative agriculture within our farming systems (minimising cultivation, maintaining good ground cover, more diverse pastures and use of deeper-rooted species) are also likely to be important for the future of pastoral farming in New Zealand (www.regenerationinternational.org). While a reduction in farmed animals is important to reduce biogenic methane production, smarter farming can actually increase farm income, even with fewer animals. We can also use native biodiversity to help address adverse effects of climate change on farms, for example through green firebreaks (Curran et al. 2018), in the provision of shade, and for erosion prone soil management.

Most importantly, pastoral farming needs to focus on adding value to products before they are exported from New Zealand. While considerable value has already been captured, more progress is and will be made by those processing farm products (e.g. milk processing companies, meat processors and timber mills). As a long-term investment in New Zealand's future, this is essential, both to enhance our economic prosperity and our environment. The increasing focus on value-added exports requires ongoing strategic thinking around supply-chains and the development of processing infrastructure able to supply value-added products. The success of the NZ Merino Company in taking an historically low value commodity (wool) and adding significant value to it, initially with fine wools but increasingly now with coarser wools, is testament to what can be achieved (www.nzmerino.co.nz). A critical part of that added value in the eyes of discerning customers and consumers is New Zealand's distinct brand story of which biodiversity and extensive pastoral farming's role in that plays a starring role. While adding value within New Zealand might not necessarily provide immediate additional return to farmers when you take into account the additional cost of getting products retail-ready, packaged, branded and shipped to multiple markets, our distinct brand story is important in the medium and longer term for being able to market our products globally to those who are prepared to pay for quality products with a good environmental story.

Tourism has been suggested as a more sustainable alternative to pastoral farming in New Zealand, but the environmental and social impacts of tourism are poorly understood and documented, and its carbon costs are not included within national-level accounting but are

known to be high (Creutzig et al. 2015). Tourism carbon costs are especially high for a country like New Zealand where access requires either long-haul flights or substantial ocean journeys for international tourists. Tourism is also sensitive to events like natural disasters such as the Christchurch and Kaikoura earthquakes, terrorist attacks, disease outbreaks (e.g. COVID-19 virus) and downturns in the global economy (as Iceland is currently experiencing). While tourism is and will continue to be an important part of the New Zealand economy, and can provide complementary income to farmers (through accommodation, farm experience and recreational opportunities), it would seem an unwise strategy to rely on tourism alone for sustaining the New Zealand economy and environment into the future. The increasing success and pressures of tourism, both on our native habitats and greenhouse gases do present significant opportunities for long-term partnerships between tourism operators and farmers seeking support for long term investment in restoring native forest that also generates income from carbon.

The ETS provides some incentive for establishing native vegetation on farms. However, the exclusion of native forests, including degraded (cutover) and regenerating native forests that were already present in 1990 from the ETS is ignoring the immense value these areas have for both carbon sequestration and biodiversity conservation. Furthermore, small (< 1 ha) post-1989 regenerating or planted native forest patches are also excluded under the ETS, yet can be vitally important for biodiversity conservation and carbon-sequestration at a regional and national scale. Providing a financial credit associated with carbon-sequestration from both pre-1990 forests and < 1 ha areas of post-1989 regenerating and planted native forests could be a critical incentive for farmers to retain and enhance biodiversity on their farms.

Rising carbon credit value can, however, also result in perverse outcomes for biodiversity (Lindenmayer et al. 2012). This is particularly relevant to rural New Zealand where regenerating native forests are under risk of being converted to exotic plantations because of their potential to earn greater carbon-credits. While landowners are unable to earn carbon credits from a post-1989 exotic plantation that was established on land that was formerly native forest, regenerating kānuka (*Kunzea* species) forest appears exempt from this. This occurs because of anomalies in the definition of ‘forest’ under the ETS where there are no carbon liabilities associated with converting pre-1990 kānuka stands to plantation species. Such perverse outcomes need to be avoided. It is critical for finding win-win outcomes for pastoral farming and biodiversity that these policy and incentive issues are addressed if we are to support a transformative landscape approach to biodiversity conservation in New Zealand. In particular, allowing farmers to earn credits for carbon-sequestration associated with pre-1990 forests and making clearance of pre-1990 kānuka subject to carbon liabilities would further incentivise retention of native woody vegetation on farms.

The cultural context for biodiversity conservation in agroecosystems is also critical (Brown et al. 2019). When farmers feel ownership for native biodiversity on their farms and it becomes an integral part of their farm and family story, they are far more likely to want to sustain and enhance it than if they are required to do this as a result of regulation. The land stewardship ethic and inter-generational thinking that characterises much of pastoral farming in New Zealand, especially sheep and beef farming (Elliott & Wakelin 2016), represents a huge opportunity for biodiversity conservation. The challenge is to incentivise, support and reward farmers for what they are currently doing and what they can do in the future, rather than taking a punitive regulatory approach towards biodiversity in agroecosystems, essentially focusing on restrictions rather than facilitation.

Supporting on-farm biodiversity management

A key factor that can support our ability to achieve the win-win outcomes advocated for in this article is the establishment of an integrated New Zealand-wide independent verification system that focuses on biodiversity outcomes and shows that farmers are actually doing the things they say they are doing (Williams et al. 2019). This is vital for being able to sell value-added farm products based on our environmental story to the world market as well as retaining the ‘social license to operate’ in New Zealand. Most international agricultural sustainability standards address biodiversity conservation, for example, including requirements for habitat protection, prohibiting clearance of certain land-cover types (including native forests), identifying priority habitat areas, and managing impacts to threatened species including measures to address invasive species (Milder et al. 2015, Englund & Berndes 2015). These standards have brought benefits for farmers through improved branding position, increased market access, an enhancement in their social licence to operate and in some cases improved prices. However, the degree to which each standard delivers environmental benefits vary (Blackman & Rivera 2011; Tayleur et al. 2017). The development of agricultural sustainability standards requires a robust evidence-based monitoring framework focused on environmental and biodiversity outcomes that are measured using objective time-series methods (Williams et al. 2019). However, such a framework needs to be developed and applied in a way that is financially viable for farmers. Stakeholders and consumers need to be confident that farmers really are doing the things they say they are doing on-farm. Ideally, the verification system will enable transparency and traceability where a consumer will be able to access information about the farm the product is sourced from and be able to obtain information on what is actually happening on that specific farm (Lernoud et al. 2017).

The verification system should be supported by farm environment planning where biodiversity (and other environmental issues) is just as central to farm management as soils, pasture and animals. To do this, a comprehensive demonstration and extension system focusing on both farmers and their advisors is required that provides information to farmers about what biodiversity is, why it is important and how it can be managed. This extension system should be independent of the regulatory system otherwise uptake will be limited (Norton & Reid 2013). Having specific biodiversity knowledge additional to what people know generally is important to support farmers. Time and money are key limitations to farmers acquiring advice on what biodiversity they have and how best to manage it, issues which such a resource can address. Development of a biodiversity support resource is critical for achieving the win-win outcome model proposed here and needs to be based on well-developed (online) extension resources, supported by independent biodiversity experts (biodiversity ambassadors *sensu* Norton & Reid 2013) who take this information out to the farming community (farmers and farm advisors).

Conclusions

To obtain win-win outcomes for biodiversity conservation, pastoral farming and New Zealanders generally, we need to avoid letting our policy systems (national, regional and district) fall back on a strict rules-based approach. This tends to stifle innovation and result in perverse outcomes. We need to reorganise our policy system to incentivise and reward farmers for good biodiversity management which includes looking after remnants of forest and wetlands, and regenerating native forest and shrubland, and improving connectivity through landscape-level integration. The approach advocated for here requires trust and partnership among all players in agroecosystems – farmers, government, food and

fibre processors, scientists, conservationists, NGOs, and the wider New Zealand population. If we redesign our policy frameworks to support and empower a true landscape scale partnership-based approach to managing and restoring biodiversity, share resources between business, public, landowners and Government, and back this with an independent verification system, then and only then will our biodiversity thrive. If we can achieve that collectively it will strongly support market access, our strong brand story to support premium primary produce, thriving tourism and most importantly a biodiverse thriving New Zealand of which all New Zealanders can be proud.

Acknowledgements

We are appreciative of comments on this article from Maurie Leyland, Richard McDowell, Nick Reid, Rhiannon Smith, David Wardle, and three anonymous reviewers. This work was funded by the Ministry of Business, Innovation and Employment (New Zealand's Biological Heritage NSC, C09X1501).

References

- Almiron N, Tafalla M 2019. Rethinking the ethical challenge in the climate deadlock: anthropocentrism, ideological denial and animal liberation. *Journal of Agricultural and Environmental Ethics* 32: 255–267.
- Balmford B, Green RE, Onial M, Phalan B, Balmford A 2019 How imperfect can land sparing be before land sharing is more favourable for wild species? *Journal of Applied Ecology* 56: 73–84.
- Beef + Lamb New Zealand 2018a. Environment strategy and implementation plan 2018-22. <https://beeflambnz.com/environment-strategy> (accessed on 28 February 2020)
- Beef + Lamb New Zealand 2018b. Compendium of New Zealand farm facts 2018. <https://beeflambnz.com/knowledge-hub/PDF/compendium-farm-facts> (accessed on 28 February 2020)
- Blackman A, Rivera J 2011. Producer-level benefits of sustainability certification. *Conservation Biology* 25: 1176–1185.
- Blaschke P, Ngapo N 2003. Review of New Zealand Environmental Farm Plans. Wellington, Ministry for the Environment. 53 p.
- Brown MA 2016. Pathways to prosperity, safeguarding biodiversity in development. Auckland, Environmental Defence Society. 74 p.
- Brown M, Kaye-Blacke B, Payne P 2019. Heartland strong: How rural New Zealand can change and thrive. Palmerston North, Massey University Press. 240 p.
- Burns BR, Barker G, Harris R, Innes J 2000. Conifers and cows: Forest survival in a New Zealand dairy landscape. In: Craig JL, Mitchell N, Saunders DA eds. *Nature conservation 5: Conservation in production environments: Managing the matrix*. Chipping Norton, Surrey Beatty & Sons. Pp. 81–89.
- Burns BR, Floyd CG, Smale MC, Arnold GC 2011. Effects of forest fragment management on vegetation condition and maintenance of canopy composition in a New Zealand pastoral landscape. *Austral Ecology* 36: 153–166.
- Butler SJ, Boccaccio L, Gregory RD, Vorisek P, Norris K 2010. Quantifying the impact of land-use change to European farmland bird populations. *Agriculture, Ecosystems & Environment* 137: 348–357.
- Campbell KL, Schotborgh HM, Wilson K-J, Ogilvie SC 2008. Diet of kereru (*Hemiphaga novaeseelandiae*) in a rural-urban landscape, Banks Peninsula, New Zealand. *Notornis* 55: 173–183.

- Creutzig F, Pochem P, Edelenbosch OY, Mattauch L, van Vuuren DP, McCollum D, Minx J 2015. Transport: A roadblock to climate mitigation? *Science* 350: 911–912.
- Crist E, Mora C, Engelman R 2017. The interaction of human population, food production, and biodiversity protection. *Science* 356: 260–264.
- Curran TJ, Perry GLW, Wyse SV, Alam MA 2018. Managing fire and biodiversity in the wildland-urban interface: A role for green firebreaks. *Fire* 2018(1): 3.
- Dairy Tomorrow 2017. Dairy tomorrow: the future of New Zealand dairying. The dairy industry strategy 2017-2025. <https://www.dairytomorrow.co.nz/> (accessed on 28 February 2020)
- Dale VH, Pearson SM, Offerman HL, O'Neill RV 1994. Relating patterns of land-use change to faunal biodiversity in the central Amazon. *Conservation Biology* 8: 1027–1036.
- de Lange P, Heenan P, Norton D., Rolfe J, Sawyer J 2010. Threatened plants of New Zealand. Canterbury University Press, Christchurch. 471 p.
- Didham RK, Barker GM, Bartlam S, Deakin EL, Denmead LH, Fisk LM, Peters JMR, Tylianakis JM, Wright HR, Schipper LA 2015. Agricultural intensification exacerbates spillover effects on soil biogeochemistry in adjacent forest remnants. *PLoS ONE* 10(1): e0116474.
- Diprose D 2018. Pressures of regulation and communities changing face of farming. Wellington, Stuff NZ. <https://www.stuff.co.nz/business/farming/102216997/pressures-of-regulation-and-communities-changing-face-of-farming> (accessed on 28 February 2020)
- Dodd M, Barker G, Burns B, Didham R, Innes J, King C, Smale M, Watts C 2011. Resilience of New Zealand indigenous forest fragments to impacts of livestock and pest mammals. *New Zealand Journal of Ecology* 35: 83–95.
- Dominati EJ, MacKay AD, Rendel JM, Smale PN 2016. Looking to the future of land evaluation and farm planning. *Journal of New Zealand Grasslands* 78: 67–72.
- Elliott MK, Wakelin RDN 2016. Drivers of top performing farmers. *Hill Country – Grassland Research and Practice Series* 16: 25–28.
- Englund O, Berndes G 2015. How do sustainability standards consider biodiversity? *Wiley Interdisciplinary Reviews: Energy and Environment* 4: 26–50.
- Ewers RM, Kliskey AD, Walker S, Rutledge D, Harding JS, Didham RK 2006. Past and future trajectories of forest loss in New Zealand. *Biological Conservation* 133: 312–325.
- Fischer J, Abson DJ, Butsic V, Chappell MJ, Ekroos J, Hanspach J, Kuemmerle T, Smith HG, von Wehrden H 2014. Land sparing versus land sharing: moving forward. *Conservation Letters* 7: 149–157.
- Foote KJ, Joy MK, Death RG 2015. New Zealand dairy farming; milking our environment for all its worth. *Environmental Management* 56: 709–720.
- Forbes AS, Wallace KJ, Buckley HL, Case BS, Clarkson BD, Norton DA 2020. Restoring mature-phase forest tree species through enrichment planting in New Zealand's lowland landscapes. *New Zealand Journal of Ecology* 44(1): 3404.
- Fukuda Y, Moller H, Burns B. 2011. Effects of organic farming, fencing and vegetation origin on spiders and beetles within shelterbelts on dairy farms. *New Zealand Journal of Agricultural Research* 54: 155-176.
- Gabzdylova B, Raffensperger JF, Castka P 2009. Sustainability in the New Zealand wine industry: drivers, stakeholders and practices. *Journal of Cleaner Production* 17: 992–998.
- Grass I, Loos J, Baensch S, Batáry P, Librán-Embíd F, Ficiciyan A, Klaus F, Riechers M, Rosa J, Tiede J, Udy K 2019. Land-sharing/-sparing connectivity landscapes for ecosystem services and biodiversity conservation. *People and Nature* 1: 262–272.

- Hutching 2019. Carbon farming can provide better returns than sheep and beef. Wellington, Stuff NZ. <https://www.stuff.co.nz/business/farming/112416878/carbon-farming-can-provide-better-returns-than-sheep-and-beef> (accessed on 28 February 2020)
- IPCC 2019a. Chapter 2: Land-Climate Interactions. IPCC, 1–186). <https://www.ipcc.ch/report/srcl/> (accessed on 28 February 2020)
- IPCC 2019b. Chapter 6: Interlinkages between desertification, land degradation, food security and GHG fluxes : synergies , trade-offs and integrated response options. <https://www.ipcc.ch/report/srcl/> (accessed on 28 February 2020)
- Kennet P, Kennett J, Johnson S, Bennion T 2010. The carbon forest: A New Zealand guide to forest carbon sinks for investors, farmers, foresters, and conservationists. Wellington, Kennett Brothers. 173 p.
- Leathwick JR 2001. New Zealand's potential forest pattern as predicted from current species-environment relationships. *New Zealand Journal of Botany* 39: 447–464.
- Leining C, Kerr S 2018. A Guide to the New Zealand Emissions Trading Scheme. Report prepared Motu Economic and Public Policy Research for the Ministry for the Environment, Wellington. 15 p.
- Lernoud J, Potts J, Sampson G, Garibay S, Lynch M, Voora V, Willer H, Wozniak J 2017. The state of sustainable markets – Statistics and emerging trends 2017. <http://www.intracen.org/publication/The-State-of-Sustainable-Markets-2018-Statistics-and-Emerging-Trends/> (accessed on 28 February 2020)
- Lindenmayer DB, Hulvey KB, Hobbs, RJ 2012. Avoiding bio-perversity from carbon sequestration solutions. *Conservation Letters* 5: 28–36.
- MacLeod CJ, Blackwell G, Moller H, Innes J, Powlesland R 2008. The forgotten 60%: bird ecology and management in New Zealand’s agricultural landscapes. *New Zealand Journal of Ecology* 32: 240–255.
- Macleod CJ, Moller H 2006. Intensification and diversification of New Zealand agriculture since 1960: An evaluation of current indicators of land use change. *Agriculture, Ecosystems and Environment* 115: 201–218.
- Manning AD, Fischer J, Lindenmayer DB 2006. Scattered trees are keystone structures – Implications for conservation. *Biological Conservation* 132: 311–321.
- Maseyk FJF, Dominati EJ, Mackay AD 2019 More than a ‘nice to have’: integrating indigenous biodiversity into agroecosystems in New Zealand. *New Zealand Journal of Ecology* 43(2): 3372.
- Maseyk FJF, Dominati EJ, White T, Mackay, AD 2017. Farmer perspectives of the on-farm and off-farm pros and cons of planted multifunctional riparian margins. *Land Use Policy* 61: 160–170.
- McArthur N, Boulton RL, Richard Y, Armstrong DP 2019. The role of pine plantations in source-sink dynamics of North Island robins. *New Zealand Journal of Ecology* 43(1): 3362.
- McDowell RW 2017. Does variable rate irrigation decrease nutrient leaching losses from grazed dairy farming? *Soil Use and Management* 33: 530–537.
- McGlone MS 2000. The origin of the indigenous grasslands of southeastern South Island in relation to pre-human woody ecosystems. *New Zealand Journal of Ecology* 25: 1–15.
- Milder JC, Arbuthnot M, Blackman A, Brooks SE, Giovannucci D, Gross L, Kennedy ET, Komives K, Lambin EF, Lee A, Meyer D, Newton P, Phalan B, Schroth G, Semroc B, Van Rikxoort H, Zrust M 2015. An agenda for assessing and improving conservation impacts of sustainability standards in tropical agriculture. *Conservation Biology* 29: 309–320.
- Ministry for the Environment 2019a. Environment Aotearoa 2019. Wellington, Ministry for the Environment. 128 p.

- Ministry for the Environment 2019b. He kura koiora I hokia: A discussion document on a proposed National Policy Statement for Indigenous Biodiversity. Wellington, Ministry for the Environment. 103 p.
- Mitchell C 2017. Farming, emissions and waste putting NZ's 'green' reputation at risk, OECD says. Wellington, Stuff NZ. <https://www.stuff.co.nz/environment/90613205/new-zealand-showing-environmental-limits-oecd-says> (accessed on 28 February 2020)
- Newbold T, Hudson LN, Hill SLL, Contu S, Lysenko I, Senior RA, Boerger L, Bennett DJ, Choimes A, Collen B, Day J, De Palma A, Diaz S, Echeverria-Londono S, Edgar MJ, Feldman A, Garon M, Harrison MLK, Alhousseini T, Ingram DJ, Itescu Y, Kattge J, Kemp V, Kirkpatrick L, Kleyer M, Correia DLP, Martin CD, Meiri S, Novosolov M, Pan Y, Phillips HRP, Purves DW, Robinson A, Simpson J, Tuck SL, Weiher E, White HJ, Ewers RM, Mace GM, Scharlemann JPW, Purvis A 2015. Global effects of land use on local terrestrial biodiversity. *Nature* 520: 45–50.
- Norton D, Reid N 2013. Nature and farming. Sustaining native biodiversity in agricultural landscapes. Melbourne, CSIRO Publishing. 29 p.
- Norton DA 1988. Managing for the long term. *Forest and Bird* 19(2): 32–34.
- Norton DA 1998. Indigenous biodiversity conservation and plantation forestry: options for the future. *New Zealand Forestry* 43(2): 34–39.
- Norton DA 2001. Is the amount and focus of ecological research in New Zealand sufficient to sustain indigenous biodiversity on private land? *New Zealand Journal of Ecology* 25: 77–82.
- Norton D, Pannell J 2018. Desk-top assessment of native vegetation on New Zealand sheep and beef farms. Report prepared for Beef+Lamb New Zealand. <https://beeflambnz.com/norton-report> (accessed on 28 February 2020)
- Norton DA, Butt J, Bergin DO 2018. Upscaling restoration of native biodiversity: A New Zealand perspective. *Ecological Management and Restoration* 19 S1: 26–35.
- Parliamentary Commissioner for the Environment 2019. Farms, forests and fossil fuels: The next great landscape transformation? Wellington, Parliamentary Commissioner for the Environment. 183 p.
- Pawson SM, Ecroyd CE, Seaton R, Shaw WB, Brockerhoff EG 2010. New Zealand's exotic plantation forests as habitats for threatened indigenous species. *New Zealand Journal of Ecology* 34: 342–355.
- Perry GLW, Wilmshurst JM, McGlone MS 2014. Ecology and long-term history of fire in New Zealand. *New Zealand Journal of Ecology* 38: 157–176.
- Poore J, Nemecek T 2018. Reducing food's environmental impacts through producers and consumers. *Science* 360: 987–992.
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE 2010. Global pollinator declines: trends, impacts and drivers. *Trends in Ecology & Evolution* 25: 345–353.
- Ramezani J, Akbaripasand A, Closs GP, Matthaei CD 2016. In-stream water quality, invertebrate and fish community health across a gradient of dairy farming prevalence in a New Zealand river catchment. *Limnologica* 61: 14–28.
- Richard Y, Armstrong DP 2010. Cost distance modelling of landscape connectivity and gap-crossing ability using radio-tracking data. *Journal of Applied Ecology* 47: 603–610.
- Ruffell J, Didham RK 2017. Conserving biodiversity in New Zealand's lowland landscapes: does forest cover or pest control have a greater effect on native birds? *New Zealand Journal of Ecology* 41: 23–33.
- Smale MC, Dodd MB, Burns BR, Power IL 2008. Long-term impacts of grazing on indigenous forest remnants on North Island hill country, New Zealand. *New Zealand Journal of Ecology* 32: 57–66.

- Tayleur C, Balmford A, Buchanan GM, Butchart SHM, Ducharme H, Green RE, Milder JC, Sanderson FJ, Thomas DHL, Vickery J, Phalan B 2017. Global coverage of agricultural sustainability standards, and their role in conserving biodiversity. *Conservation Letters* 10: 610–618.
- Te Uru Rākau 2018. Guide to One Billion Trees.
<https://www.teururakau.govt.nz/dmsdocument/31860-guide-to-the-one-billion-trees-programme-2018> (accessed on 28 February 2020)
- Tilman D 1999. Global environmental impacts of agricultural expansion: The need for sustainable and efficient practices. *Proceedings of the National Academy of Sciences* 96: 5995–6000.
- Timar L. 2016. Does money grow on trees? Mitigation under climate policy in a heterogeneous sheep-beef sector (No. 16_09).
https://ideas.repec.org/p/mtu/wpaper/16_09.html (accessed on 28 February 2020)
- Timmins SM, Williams PA 1991. Weed numbers in New Zealand's forest and scrub reserves. *New Zealand Journal of Ecology* 15: 153–162.
- Wallace N 2019. Tree growth worries meat firms. *Farmers Weekly*.
<https://farmersweekly.co.nz/section/agribusiness/view/tree-growth-worries-meat-firms> (accessed on 28 February 2020)
- Williams J, Smith R, Ball A, Reid N, Kahn L 2019. Harmonisation of on-farm metrics for sustainability assessment of Australian agricultural industries. *Farm Policy Journal*.
- Wilmshurst JM, Hunt TL, Lipo CP, Anderson AJ 2011. High-precision radiocarbon dating shows recent and rapid initial human colonization of East Polynesia. *Proceedings of the National Academy of Sciences* 108: 1815–1820.

Received 20 August 2019; accepted 3 February 2020
Editorial board member: David Wardle